

## POSTER PRESENTATION

## Open Access

# Bacterial colonization of central venous catheters after heart valve surgery: a risk factor study

A De La Chapelle<sup>1\*</sup>, A Mihoubi<sup>2</sup>, N Balarac<sup>3</sup>, M Raucoules-Aime<sup>4</sup>, P Dellamonica<sup>5</sup>

From 3rd International Conference on Prevention and Infection Control (ICPIC 2015)  
Geneva, Switzerland. 16-19 June 2015

## Introduction

In the presence of a prosthetic heart valve, the colonization of a central venous catheter (CVC) has been implicated as a risk factor for endocarditis. Avoiding bacterial colonization of Central Venous Catheters (CVCs) is an everyday challenge for clinicians

## Objectives

This study evaluated EuroSCORE (Score use for predicting surgical mortality) and the preoperative presence of diabetes to identify patients at a higher risk of a bacterial contamination of the central venous catheter (CVC) after heart valve replacement or valvuloplasty.

## Methods

An observational evaluation was conducted from January 2006 through January 2013 on prospective data submitted to a database. 1324 consecutive patients after valve surgery were included (from January 2006 to December 2010). The systematic cultures of CVC were performed for all patients regardless of infection symptoms. A long-term monitoring (2 years) was done by phone call (up to January 2013). Patients suspected of prosthetic valve endocarditis had an echocardiography, and a blood analysis (particularly, blood cultures). Statistical analysis was processed by Systat 11 using Chi 2 test; Student's t test; Kruskal-Wallis test; Fisher exact test and a 'step-by-step' logistic regression. A p value < 0.05 was considered statistically significant.

## Results

The catheter-related bacteraemia was 5‰ or 0.84/1000 catheter days. The values of the additive and logistic EuroSCOREs were not significantly higher in case of CVC colonization. EuroSCOREs higher than 6 were significantly but only moderately involved with the

occurrence of CVC colonization (p = 0.034) (Odds Ratio 1.76; 95% CI [1.04; 2.97]). Diabetes was significantly but moderately associated with CVC colonization (p = 0.041) (Odds Ratio 1.87; 95% CI [1.02; 3.43]). Among parameters of euroSCORE, an ejection fraction of < 30% was closely related to CVC colonization (p = 0.004) (Odds Ratio 4.91; 95% CI [1.65; 14.55]). The catheter's exposure influenced significantly catheter colonization.

## Conclusion

A poor left ventricular function is the main risk factor in bacterial colonization of CVC. Global risk score and preexisting diabetes are not useful to predict postoperative CVC colonization

## Disclosure of interest

None declared.

## Authors' details

<sup>1</sup>Anesthesiology, institut Arnault Tzanck, Saint Laurent du Var, France.

<sup>2</sup>Cardiology, institut Arnault Tzanck, France. <sup>3</sup>Biostatistic, institut Arnault Tzanck, France. <sup>4</sup>Anesthesiology, CHU, Nice, France. <sup>5</sup>Infectious diseases, CHU, Nice, France.

Published: 16 June 2015

doi:10.1186/2047-2994-4-S1-P209

**Cite this article as:** De La Chapelle et al.: Bacterial colonization of central venous catheters after heart valve surgery: a risk factor study. *Antimicrobial Resistance and Infection Control* 2015 **4**(Suppl 1):P209.

<sup>1</sup>Anesthesiology, institut Arnault Tzanck, Saint Laurent du Var, France  
Full list of author information is available at the end of the article